



PCT/GB 00/03760


#2



INVESTOR IN PEOPLE

6B 00/3760

4




The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

Dated 6 October 2000

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

The
Patent
Office



RECEIVED: 1 OCT 1999
10.10.1999

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form))

The Patent Office

Cardiff Road
Newport
Gwent NP9 1RH

1. Your reference	P.77933 JCI		
2. Patent application number (The Patent Office will fill in this part)	01 OCT 1999	9923306.6	
3. Full name, address and postcode of the or of each applicant (underline all surnames)	ISIS INNOVATION LIMITED 2 South Parks Road, Oxford OX1 3UB United Kingdom		
Patents ADP number (if you know it)	399 856 4001		
If the applicant is a corporate body, give the country/state of its incorporation	GB		
4. Title of the invention	DIAGNOSTIC AND THERAPEUTIC EPITOPE, AND TRANSGENIC PLANT		
5. Name of your agent (if you have one)	J A KEMP & CO		
"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)	14 SOUTH SQUARE GRAY'S INN LONDON WC1R 5LX		
Patents ADP number (if you know it)	26001		
6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	Country	Priority application number (if you know it)	Date of filing (day / month / year)
7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier application		Date of filing (day / month / year)
8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer "Yes" if: a) any applicant named in part 3 is not an inventor, or b) there is an inventor who is not named as an applicant, or c) any named applicant is a corporate body: See note (d))	YES		

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description

42

Claim(s)

9

Abstract

1

Drawing(s)

11

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents
(please specify)

11. I/We request the grant of a patent on the basis of this application

Signature

J. A. V. ...

Date 1 October 1999

12. Name and daytime telephone number of person to contact in the United Kingdom
- J.C. IRVINE
0171 405 3292

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

a) If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.

b) Write your answers in capital letters using black ink or you may type them.

c) If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.

d) If you have answered "Yes" Patents Form 7/77 will need to be filed.

e) Once you have filled in the form you must remember to sign and date it.

f) For details of the fee and ways to pay please contact the Patent Office

DIAGNOSTIC AND THERAPEUTIC EPITOPE, AND TRANSGENIC PLANT

The invention relates to the diagnosis and therapy of coeliac disease, and to a gliadin protein which does not cause coeliac disease.

5 An immune reaction to gliadin (a component of gluten) in the diet causes coeliac disease. It is known that immune responses in the intestinal tissue preferentially respond to gliadin which has been modified by an intestinal transglutaminase. Coeliac disease is diagnosed by detection of anti-endomysial antibodies, but this requires confirmation by the finding of a lymphocytic
10 inflammation in intestinal biopsies. The taking of such a biopsy is inconvenient for the patient.

Investigators have previously assumed that only intestinal T cell responses provide an accurate indication of the immune response against gliadins. Therefore they have concentrated on the investigation of T cell responses in intestinal tissue¹.
15 Gliadin epitopes which require transglutaminase modification (before they are recognised by the immune system) are known².

The inventors have found the immunodominant T cell epitope recognised by the immune system in coeliac disease, and have shown that this is recognised by T cells in the peripheral blood of individuals with coeliac disease. Such T cells were
20 found to be present at high enough frequencies to be detectable without restimulation (i.e. a 'fresh response' detection system could be used). The epitope was identified using a non-T cell cloning based method which provided a more accurate reflection of the epitopes being recognised. The immunodominant epitope requires transglutaminase modification (causing substitution of a particular glutamine to
25 glutamate) before immune system recognition.

Based on this work the inventors have developed a test which can be used to diagnose coeliac disease at an early stage. The test may be carried out on a sample from peripheral blood and therefore an intestinal biopsy is not required. The test is more sensitive than the antibody tests which are currently being used.

30 The invention thus provides a method of diagnosing coeliac disease, or susceptibility to coeliac disease, in an individual comprising:

(a) contacting a sample from the host with an agent selected from (i) the epitope comprising sequence which is: SEQ ID NO:1 or 2, or an equivalent sequence from a naturally occurring homologue of the gliadin represented by SEQ ID NO:3, (ii) an epitope comprising sequence comprising: SEQ ID NO:1, or an equivalent sequence from a naturally occurring homologue of the gliadin represented by SEQ ID NO:3, which epitope is an isolated oligopeptide derived from a gliadin protein, (iii) an analogue of (i) or (ii) which is capable of being recognised by a T cell receptor that recognises (i) or (ii), which in the case of a peptide analogue is not more than 50 amino acids in length, or (iv) a product comprising two or more agents as defined in (i), (ii) or (iii), and (b) determining *in vitro* whether T cells in the sample recognise the agent, recognition by the T cells indicating that the individual has, or is susceptible to, coeliac disease.

The invention also provides use of the agent for the preparation of a diagnostic means for use in a method of diagnosing coeliac disease, or susceptibility to coeliac disease, in an individual, said method comprising determining whether T cells of the individual recognise the agent, recognition by the T cells indicating that the individual has, or is susceptible to, coeliac disease.

The finding of an immunodominant epitope which is modified by transglutaminase also allows diagnosis of coeliac disease based on determining whether other types of immune response to this epitope are present. Thus the invention also provides a method of diagnosing coeliac disease, or susceptibility to coeliac disease, in an individual comprising determining the presence of an antibody that binds to the epitope in a sample from the individual, the presence of the antibody indicating that the individual has, or is susceptible to, coeliac disease.

The invention additionally provides the agent, optionally in association with a carrier, for use in a method of treating or preventing coeliac disease by tolerising T cells which recognise the agent. Also provided is an antagonist of a T cell which has a T cell receptor that recognises (i) or (ii), optionally in association with a carrier, for use in a method of treating or preventing coeliac disease by antagonising such T cells. Additionally provided is the agent or an analogue that binds an antibody (that binds the agent) for use in a method of treating or preventing coeliac disease in an

individual by tolerising the individual to prevent the production of such an antibody.

The invention provides a method of determining whether a composition is capable of causing coeliac disease comprising determining whether a protein capable of being modified by a transglutaminase to an oligopeptide sequence as defined
5 above is present in the composition, the presence of the protein indicating that the composition is capable of causing coeliac disease.

The invention also provides a mutant gliadin protein whose wild-type sequence can be modified by a transglutaminase to a sequence that comprises an epitope comprising sequence as defined above, but which mutant gliadin protein has
10 been modified in such a way that it does not contain sequence which can be modified by a transglutaminase to a sequence that comprises such an epitope comprising sequence; or a fragment of such a mutant gliadin protein which is at least 15 amino acids long and which comprises sequence which has been modified in said way.

Additionally the invention provides a protein that comprises a sequence
15 which is able to bind to a T cell receptor, which T cell receptor recognises the agent, and which sequence is able to cause antagonism of a T cell that carries such a T cell receptor.

Additionally the invention provides a food that comprises the proteins defined above.

20 The invention is illustrated by the accompanying drawings in which:

Figure 1 shows freshly isolated PBMC (peripheral blood mononuclear cell) IFN γ ELISPOT responses (vertical axis shows spot forming cells per 10⁶ PBMC) to transglutaminase (tTG)-treated and untreated peptide pool 3 (each peptide 10 μ g/ml) including five overlapping 15mers spanning A-gliadin 51-85 (see Table 1) and a-
25 chymotrypsin-digested gliadin (40 μ g/ml) in Coeliac disease Subject 1, initially in remission following a gluten free diet then challenged with 200g bread daily for three days from day 1 (a). PBMC IFN γ ELISPOT responses by Subject 2 to tTG-treated A-gliadin peptide pools 1-10 spanning the complete A-gliadin protein during ten day bread challenge (b). The horizontal axis shows days after commencing bread.

30 Figure 2 shows PBMC IFN γ ELISPOT responses to tTG-treated peptide pool 3 (spanning A-gliadin 51-85) in 7 individual Coeliac disease subjects (vertical axis

shows spot forming cells per 10^6 PBMC), initially in remission on gluten free diet, challenged with bread for three days (days 1 to 3). The horizontal axis shows days after commencing bread. (a). PBMC IFN γ Elispot responses to tTG-treated overlapping 15mer peptides included in pool 3; bars represent the mean (\pm SEM) response to individual peptides (10 μ g/ml) in 6 Coeliac disease subjects on day 6 or 7 (b). (In individual subjects, ELISPOT responses to peptides were calculated as a % of response elicited by peptide 12 - as shown by the vertical axis.)

Figure 3 shows PBMC IFN γ ELISPOT responses to tTG-treated truncations of A-gliadin 56-75 (0.1 μ M). Bars represent the mean (\pm SEM) in 5 Coeliac disease subjects. (In individual subjects, responses were calculated as the % of the maximal response elicited by any of the peptides tested.)

Figure 4 shows how the minimal structure of the dominant A-gliadin epitope was mapped using tTG-treated 7-17mer A-gliadin peptides (0.1 μ M) including the sequence, PQQQLPY (A-gliadin 62-68) (a), and the same peptides without tTG treatment but with the substitution Q \rightarrow E65 (b). Each line represents PBMC IFN γ ELISPOT responses in each of three Coeliac disease subjects on day 6 or 7 after bread was ingested on days 1-3. (In individual subjects, ELISPOT responses were calculated as a % of the response elicited by the 17mer, A-gliadin 57-73.)

Figure 5 shows the amino acids which were deamidated by tTG. A-gliadin 56-75 (LQLQPFPPQQLPYQPQSFP) (0.1 μ M) was incubated with tTG (50 μ g/ml) at 37°C for 2 hours. A single product was identified and purified by reverse phase HPLC. Amino acid analysis allowed % deamidation (Q \rightarrow E) of each Gln residue in A-gliadin 56-75 attributable to tTG to be calculated (vertical axis).

Figure 6 shows the effect of substituting Q \rightarrow E in A-gliadin 57-73 at other positions in addition to Q65 using the 17mers: QLQPFPPQPELPYPQPES (E57,65), QLQPFPPQPELPYPQPES (E65,72), ELQPFPPQPELPYPQPES (E57, 65, 72), and QLQPFPPQPELPYPQPQS (E65) in three Coeliac disease subjects on day 6 or 7 after bread was ingested on days 1-3. Vertical axis shows % of the E65 response.

Figure 7 shows that tTG treated A-gliadin 56-75 (0.1 μ M) elicited IFN-g ELISPOT responses in (a) CD4 and CD8 magnetic bead depleted PBMC. (Bars represent CD4 depleted PBMC responses as a % of CD8 depleted PBMC responses;

spot forming cells per million CD8 depleted PBMC were: Subject 4: 29, and Subject 6: 535). (b) PBMC IFN γ ELISPOT responses (spot forming cells/million PBMC) after incubation with monoclonal antibodies to HLA-DR (L243), -DQ (L2) and -DP (B7.21) (10 μ g/ml) 1h prior to tTG-treated 56-75 (0.1 μ M) in two Coeliac disease subjects homozygous for HLA-DQ a1*0501, b1*0201.

Figure 8 shows the effect of substituting Glu at position 65 for other amino acids in the immunodominant epitope. The vertical axis shows the % response in the 3 subjects in relation to the immunodominant epitope.

Figure 9 shows the immunoreactivity of naturally occurring gliadin peptides (measuring responses from 3 subjects) which contain the sequence PQLPY with (shaded) and without (clear) transglutaminase treatment.

Detailed description of the invention

The term 'coeliac disease' encompasses a spectrum of conditions caused by varying degrees of gluten sensitivity, including a severe form characterised by a flat small intestinal mucosa (hyperplastic villous atrophy) and other forms characterised by milder symptoms.

The individual mentioned above (in the context of diagnosis or therapy) is human. They may have coeliac disease (symptomatic or asymptomatic) or be suspected of having it. They may be on a gluten free diet. They may be in an acute phase response (for example they may have coeliac disease, but have only ingested gluten in the last 24 hours before which they had been on a gluten free diet for 14 to 28 days).

The individual may be susceptible to coeliac disease, such as a genetic susceptibility (determined for example by the individual having relatives with coeliac disease or possessing genes which cause predisposition to coeliac disease).

The agent

The agent is typically a peptide, for example of length 7 to 50 amino acids, such as 10 to 40, or 15 to 30 amino acids in length.

SEQ ID NO:1 is PQPELPY. SEQ ID NO:2 is QLQPFPPQPELPYPQPQS.

SEQ ID NO:3 is shown in Table 1 and is the sequence of a whole A-gliadin. The glutamate at position 4 of SEQ ID NO:1 (equivalent to position 9 of SEQ ID NO:2) is generated by transglutaminase treatment of A-gliadin.

The agent may be the peptide represented by SEQ ID NO:1 or 2 or an epitope comprising sequence that comprises SEQ ID NO:1 which is an isolated oligopeptide derived from a gliadin protein; or an equivalent of these sequences from a naturally occurring gliadin protein which is a homologue of SEQ ID NO:3. Thus the epitope may be a derivative of the protein represented by SEQ ID NO:3. Such a derivative is typically a fragment of the gliadin, or a mutated derivative of the whole protein or fragment. Therefore the epitope of the invention does not include this naturally occurring whole gliadin protein, and does not include other whole naturally occurring gliadins.

The epitope may thus be a fragment of A-gliadin (e.g. SEQ ID NO:3), which comprises the sequence of SEQ ID NO:1, obtainable by treating (fully or partially) with transglutaminase, i.e. with 1, 2, 3 or more glutamines substituted to glutamates (including the substitution within SEQ ID NO:1).

Such fragments may be or may include the sequences represented by positions 55 to 70, 58 to 73, 61 to 77 of SEQ ID NO:3 shown in Table 1. Typically such fragments will be recognised by T cells to at least the same extent that the peptides represented by SEQ ID NO:1 or 2 are recognised in any of the assays described herein using samples from coeliac disease patients.

In the case where the epitope comprises a sequence equivalent to the above epitopes (including fragments) from another gliadin protein (e.g. any of the gliadin proteins mentioned herein or any gliadins which cause coeliac disease), such equivalent sequences will correspond to a fragment of a gliadin protein typically treated (partially or fully) with transglutaminase. Such equivalent peptides can be determined by aligning the sequences of other gliadin proteins with SEQ ID NO:3 (for example using any of the programs mentioned herein). Transglutaminase is commercially available (e.g. Sigma T-5398). Table 4 provides examples of suitable equivalent sequences.

The agent which is an analogue is capable of being recognised by a TCR

which recognises (i) or (ii). Therefore generally when the analogue is added to T cells in the presence of (i) or (ii), typically also in the presence of an antigen presenting cell (APC) (such as any of the APCs mentioned herein), the analogue inhibits the recognition of (i) or (ii), i.e. the analogue is able to compete with (i) or (ii) in such a system.

The analogue may be one which is capable of binding the TCR which recognises (i) or (ii). Such binding can be tested by standard techniques. Such TCRs can be isolated from T cells which have been shown to recognise (i) or (ii) (e.g. using the method of the invention). Demonstration of the binding of the analogue to the TCRs can then shown by determining whether the TCRs inhibit the binding of the analogue to a substance that binds the analogue, e.g. an antibody to the analogue. Typically the analogue is bound to a class II MHC molecule (e.g. HLA-DQ2) in such an inhibition of binding assay.

Typically the analogue inhibits the binding of (i) or (ii) to a TCR. In this case the amount of (i) or (ii) which can bind the TCR in the presence of the analogue is decreased. This is because the analogue is able to bind the TCR and therefore competes with (i) or (ii) for binding to the TCR.

T cells for use in the above binding experiments can be isolated from patients with coeliac disease, for example with the aid of the method of the invention. Other binding characteristics of the analogue may also be the same as (i) or (ii), and thus typically the analogue binds to the same MHC class II molecule to which the peptide binds (HLA-DQ2). The analogue typically binds to antibodies specific for (i) or (ii), and thus inhibits binding of (i) or (ii) to such antibodies.

The analogue is typically a peptide. It may have homology with (i) or (ii), typically at least 70% homology, preferably at least 80, 90%, 95%, 97% or 99% homology with (i) or (ii), for example over a region of at least 15 more (such as the entire length of the analogue and/or (i) or (ii), or across the region which contacts the TCR or binds the MHC molecule) contiguous amino acids. Methods of measuring protein homology are well known in the art and it will be understood by those of skill in the art that in the present context, homology is calculated on the basis of amino acid identity (sometimes referred to as "hard homology").

For example the UWGCG Package provides the BESTFIT program which can be used to calculate homology (for example used on its default settings) (Devereux *et al* (1984) *Nucleic Acids Research* 12, p387-395). The PILEUP and BLAST algorithms can be used to calculate homology or line up sequences (typically on their default settings), for example as described in Altschul S. F. (1993) *J Mol Evol* 36:290-300; Altschul, S, F *et al* (1990) *J Mol Biol* 215:403-10.

Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pair (HSPs) by identifying short words of length W in the query sequence that either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighbourhood word score threshold (Altschul *et al*, supra). These initial neighbourhood word hits act as seeds for initiating searches to find HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Extensions for the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLAST program uses as defaults a word length (W) of 11, the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1992) *Proc. Natl. Acad. Sci. USA* 89: 10915-10919) alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.

The BLAST algorithm performs a statistical analysis of the similarity between two sequences; see e.g., Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90: 5873-5787. One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a sequence is considered similar to another sequence if the smallest sum probability in comparison of the first sequence to the second

sequence is less than about 1, preferably less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

The homologous peptide analogues typically differ from (i) or (ii) by 1, 2, 3, 4, 5, 6, 7, 8 or more mutations (which may be substitutions, deletions or insertions).

5 These mutation may be measured across any of the regions mentioned above in relation to calculating homology. The substitutions are preferably 'conservative'.

These are defined according to the following Table. Amino acids in the same block in the second column and preferably in the same line in the third column may be substituted for each other:

10

ALIPHATIC	Non-polar	G A P
		I L V
	Polar - uncharged	C S T M
		N Q
	Polar - charged	D E
		K R
AROMATIC		H F W Y

15

Typically the amino acids in the analogue at the equivalent positions to amino acids in (i) or (ii) which contribute to binding the MHC molecule or are responsible for the recognition by the TCR, are the same or are conserved.

20

Typically the analogue peptide comprises one or more modifications, which may be natural post-translation modifications or artificial modifications. The modification may provide a chemical moiety (typically by substitution of a hydrogen, e.g. of a C-H bond), such as an amino, acetyl, hydroxy or halogen (e.g. fluorine) group or carbohydrate group. Typically the modification is present on the N or C terminus.

25

The analogue may comprise one or more non-natural amino acids, for example amino acids with a side chain different from natural amino acids. Generally, the non-natural amino acid will have an N terminus and/or a C terminus. The non-natural amino acid may be an L- or a D- amino acid.

The analogue typically has a shape, size, flexibility or electronic configuration which is substantially similar to (i) or (ii). It is typically a derivative of (i) or (ii). In one embodiment the analogue is a fusion protein comprising the sequence of SEQ ID NO:1 or 2, or any of the other peptides mentioned herein; and
5 non-gliadin sequence.

In one embodiment the analogue is or mimics (i) or (ii) bound to a MHC class II molecule. 2, 3, 4 or more of such complexes may be associated or bound to each other, for example using a biotin/streptavidin based system, in which typically 2, 3 or 4 biotin labelled MHC molecules bind to a streptavidin moiety. This analogue
10 typically inhibits the binding of the (i) or (ii)/MHC Class II complex to a TCR or antibody which is specific for the complex.

The analogue is typically an antibody or a fragment of an antibody, such as a Fab or (Fab)₂ fragment. The analogue may be immobilised on a solid support, particularly an analogue which mimics peptide bound to a MHC molecule.

15 The analogue is typically designed by computational means and then synthesised using methods known in the art. Alternatively the analogue can be selected from a library of compounds. The library may be a combinatorial library or a display library, such as a phage display library. The library of compounds may be expressed in the display library in the form of being bound to a MHC class II
20 molecule, such as HLA-DQ2. Analogues are generally selected from the library based on their ability to mimic the binding characteristics (i) or (ii). Thus they may be selected based on ability to bind a TCR or antibody which recognises (i) or (ii).

Typically analogues will be recognised by T cells to at least the same extent as any of the agents (i) or (ii), for example at least to the same extent as the
25 equivalent epitope and preferably to the same extent as the peptide represented by SEQ ID NO:2, is recognised in any of the assays described herein, typically using T cells from coeliac disease patients. Analogues may be recognised to these extents *in vivo* and thus may be able to induce coeliac disease symptoms to at least the same extent as any of the agents mentioned herein (e.g. in a human patient or animal
30 model).

Analogues may be identified in a method comprising determining whether a

candidate substance is recognised by a T cell receptor that recognises an epitope of the invention, recognition of the substance indicating that the substance is an analogue. Such TCRs may be any of the TCRs mentioned herein, and may be present on T cells. Any suitable assay mentioned herein can be used to identify the analogue. In one embodiment this method is carried out *in vivo*. As mentioned above preferred analogues are recognised to at least the same extent as the peptide SEQ ID NO:2, and so the method may be used to identify analogues which are recognised to this extent.

In one embodiment the method comprises determining whether a candidate substance is able to inhibit the recognition of an epitope of the invention, inhibition of recognition indicating that the substance is an analogue.

The agent may be a product comprising at least 2, 5, 10 or 20 agents as defined by (i), (ii) or (iii). Typically the composition comprises epitopes of the invention (or equivalent analogues) from different gliadins, such as any of the species or variety of or types of gliadin mentioned herein. Preferred compositions comprise at least one epitope of the invention, or equivalent analogue, from all of the gliadins present in any of the species or variety mentioned herein, or from 2, 3, 4 or more of the species mentioned herein (such as from the panel of species consisting of wheat, rye, barley, oats and triticale).

Diagnosis

As mentioned above the method of diagnosis of the invention may be based on the detection of T cells which bind the agent or on the detection of antibodies that recognise the agent.

The T cells which recognise the agent in the method (which includes the use mentioned above) are generally T cells which have been pre-sensitised *in vivo* to gliadin. As mentioned above such antigen-experienced T cells have been found to be present in the peripheral blood.

In the method the T cells can be contacted with the agent *in vitro* or *in vivo*, and determining whether the T cells recognise the agent can be performed *in vitro* or *in vivo*. Thus the invention provides the agent for use in a method of diagnosis

practiced on the human body. Different agents are provided for simultaneous, separate or sequential use in such a method.

The *in vitro* method is typically carried out in aqueous solution into which the agent is added. The solution will also comprise the T cells (and in certain
5 embodiments the APCs discussed below). The term 'contacting' as used herein includes adding the particular substance to the solution.

Determination of whether the T cells recognise the agent is generally done by detecting a change in the state of the T cells in the presence of the agent or determining whether the T cells bind the agent. The change in state is generally
10 caused by antigen specific functional activity of the T cell after the TCR binds the agent. The change of state may be measured inside (e.g. change in intracellular expression of proteins) or outside (e.g. detection of secreted substances) the T cells.

The change in state of the T cell may be the start of or increase in secretion of a substance from the T cell, such as a cytokine, especially IFN- γ , IL-2 or TNF- α .

15 Determination of IFN- γ secretion is particularly preferred. The substance can typically be detected by allowing it to bind to a specific binding agent and then measuring the presence of the specific binding agent/substance complex. The specific binding agent is typically an antibody, such as polyclonal or monoclonal antibodies. Antibodies to cytokines are commercially available, or can be made
20 using standard techniques.

Typically the specific binding agent is immobilised on a solid support. After the substance is allowed to bind the solid support can optionally be washed to remove material which is not specifically bound to the agent. The agent/substance complex may be detected by using a second binding agent which will bind the
25 complex. Typically the second agent binds the substance at a site which is different from the site which binds the first agent. The second agent is preferably an antibody and is labelled directly or indirectly by a detectable label.

Thus the second agent may be detected by a third agent which is typically labelled directly or indirectly by a detectable label. For example the second agent
30 may comprise a biotin moiety, allowing detection by a third agent which comprises a streptavidin moiety and typically alkaline phosphatase as a detectable label.

In one embodiment the detection system which is used is the *ex-vivo* ELISPOT assay described in WO 98/23960. In that assay IFN- γ secreted from the T cell is bound by a first IFN- γ specific antibody which is immobilised on a solid support. The bound IFN- γ is then detected using a second IFN- γ specific antibody which is labelled with a detectable label. Such a labelled antibody can be obtained from MABTECH (Stockholm, Sweden). Other detectable labels which can be used are discussed below.

The change in state of the T cell which can be measured may be the increase in the uptake of substances by the T cell, such as the uptake of thymidine. The change in state may be an increase in the size of the T cells, or proliferation of the T cells, or a change in cell surface markers on the T cell.

In one embodiment the change of state is detected by measuring the change in the intracellular expression of proteins, for example the increase in intracellular expression of any of the cytokines mentioned above. Such intracellular changes may be detected by contacting the inside of the T cell with a moiety that binds the expressed proteins in a specific manner and which allows sorting of the T cells by flow cytometry.

In one embodiment when binding the TCR the agent is bound to an MHC class II molecule (typically HLA-DQ2), which is typically present on the surface of an antigen presenting cell (APC). However as mentioned herein other agents can bind a TCR without the need to also bind an MHC molecule.

Generally the T cells which are contacted in the method are taken from the individual in a blood sample, although other types of samples which contain T cells can be used. The sample may be added directly to the assay or may be processed first. Typically the processing may comprise diluting of the sample, for example with water or buffer. Typically the sample is diluted from 1.5 to 100 fold, for example 2 to 50 or 5 to 10 fold.

The processing may comprise separation of components of the sample. Typically mononuclear cells (MCs) are separated from the samples. The MCs will comprise the T cells and APCs. Thus in the method the APCs present in the separated MCs can present the peptide to the T cells. In another embodiment only T

cells, such as only CD4 T cells, can be purified from the sample. PBMCs, MCs and T cells can be separated from the sample using techniques known in the art, such as those described in Lalvani *et al* (1997) *J.Exp. Med.* 186, p859-865.

In one embodiment the T cells used in the assay are in the form of unprocessed or diluted samples, or are freshly isolated T cells (such as in the form of freshly isolated MCs or PBMCs) which are used directly *ex vivo*, i.e. they are not cultured before being used in the method. Thus the T cells have not been restimulated in an antigen specific manner *in vitro*. However the T cells can be cultured before use, for example in the presence of one or more of the agents, and generally also exogenous growth promoting cytokines. During culturing the agent(s) are typically present on the surface of APCs, such as the APC used in the method. Pre-culturing of the T cells may lead to an increase in the sensitivity of the method. Thus the T cells can be converted into cell lines, such as short term cell lines (for example as described in Ota *et al* (1990) *Nature* 346, p183-187).

The APC which is typically present in the method may be from the same individual as the T cell or from a different host. The APC may be a naturally occurring APC or an artificial APC. The APC is a cell which is capable of presenting the peptide to a T cell. It is typically a B cell, dendritic cell or macrophage. It is typically separated from the same sample as the T cell and is typically co-purified with the T cell. Thus the APC may be present in MCs or PBMCs. The APC is typically a freshly isolated *ex vivo* cell or a cultured cell. It may be in the form of a cell line, such as a short term or immortalised cell line. The APC may express empty MHC class II molecules on its surface.

In the method one or more (different) agents may be used. Typically the T cells derived from the sample can be placed into an assay with all the agents which it is intended to test or the T cells can be divided and placed into separate assays each of which contain one or more of the agents.

The invention also provides the agents such as two or more of any of the agents mentioned herein (e.g. the combinations of agents which are present in the composition agent discussed above) for simultaneous separate or sequential use (eg. for *in vivo* use).

In one embodiment agent *per se* is added directly to an assay comprising T cells and APCs. As discussed above the T cells and APCs in such an assay could be in the form of MCs. When agents which can be recognised by the T cell without the need for presentation by APCs are used then APCs are not required. Analogues which mimic the original (i) or (ii) bound to a MHC molecule are an example of such an agent.

In one embodiment the agent is provided to the APC in the absence of the T cell. The APC is then provided to the T cell, typically after being allowed to present the agent on its surface. The peptide may have been taken up inside the APC and presented, or simply be taken up onto the surface without entering inside the APC.

The duration for which the agent is contacted with the T cells will vary depending on the method used for determining recognition of the peptide. Typically 10^5 to 10^7 , preferably 5×10^5 to 10^6 PBMCs are added to each assay. In the case where agent is added directly to the assay its concentration is from 10^{-1} to $10^3 \mu\text{g/ml}$, preferably 0.5 to $50 \mu\text{g/ml}$ or 1 to $10 \mu\text{g/ml}$.

Typically the length of time for which the T cells are incubated with the agent is from 4 to 24 hours, preferably 6 to 16 hours. When using *ex vivo* PBMCs it has been found that 0.3×10^6 PBMCs can be incubated in $10 \mu\text{g/ml}$ of peptide for 12 hours at 37°C .

The determination of the recognition of the agent by the T cells may be done by measuring the binding of the agent to the T cells (this can be carried out using any suitable binding assay format discussed herein). Typically T cells which bind the agent can be sorted based on this binding, for example using a FACS machine. The presence of T cells which recognise the agent will be deemed to occur if the frequency of cells sorted using the agent is above a 'control' value. The frequency of antigen-experienced T cells is generally 1 in 10^6 to 1 in 10^3 , and therefore whether or not the sorted cells are antigen-experienced T cells can be determined.

The determination of the recognition of the agent by the T cells may be measured *in vivo*. Typically the agent is administered to the host and then a response which indicates recognition of the agent may be measured. The agent is typically administered intradermally or epidermally. The agent is typically administered by

contacting with the outside of the skin, and may be retained at the site with the aid of a plaster or dressing. Alternatively the agent may be administered by needle, such as by injection, but can also be administered by other methods such as ballistics (e.g. the ballistics techniques which have been used to deliver nucleic acids). EP-A-0693119 describes techniques which can typically be used to administer the agent. Typically from 0.001 to 1000 µg, for example from 0.01 to 100 µg or 0.1 to 10 µg of agent is administered.

In one embodiment a product can be administered which is capable of providing the agent *in vivo*. Thus a polynucleotide capable of expressing the agent can be administered, typically in any of the ways described above for the administration of the agent. The polynucleotide typically has any of the characteristics of the polynucleotide provided by the invention which is discussed below. The agent is expressed from the polynucleotide *in vivo*. Typically from 0.001 to 1000 µg, for example from 0.01 to 100 µg or 0.1 to 10 µg of polynucleotide is administered.

Recognition of the agent administered to the skin is typically indicated by the occurrence of inflammation (e.g. induration, erythema or oedema) at the site of administration. This is generally measured by visual examination of the site.

The method of diagnosis based on the detection of an antibody that binds the agent is typically carried out by contacting a sample from the individual (such as any of the samples mentioned here, optionally processed in any manner mentioned herein) with the agent and determining whether an antibody in the sample binds the agent, such a binding indicating that the individual has, or is susceptible to coeliac disease. Any suitable format of binding assay may be used, such as any such format mentioned herein.

Therapy

The identification of the immunodominant epitope allows the therapeutic products to be made which target the T cells which recognise this epitope (such T cells being ones which participate in the immune response against gliadin). This finding also allows the prevention or treatment of coeliac disease by suppressing (by

tolerisation) an antibody or T cell response to the epitope.

Certain agents of the invention bind the TCR which recognises the epitope of the invention (as measured using any of the binding assays discussed above) and cause tolerisation of the T cell that carries the TCR. Such agents, optionally in association with a carrier, can therefore be used to prevent or treat coeliac disease.

Generally tolerisation can be caused by the same peptides which can (after being recognised by the TCR) cause antigen specific functional activity of the T cell (such as any such activity mentioned herein, e.g. secretion of cytokines). Such agents cause tolerisation when they are presented to the immune system in a 'tolerising' context.

Tolerisation leads to a decrease in the recognition of a T cell or antibody epitope by the immune system. In the case of a T cell epitope this can be caused by the deletion or anergising of T cells which recognise the epitope. Thus T cell activity (for example as measured in suitable assays mentioned herein) in response to the epitope is decreased. Tolerisation of an antibody response means that a decreased amount of specific antibody to the epitope is produced when the epitope is administered.

Methods of presenting antigens to the immune system in such a context are known and are described for example in Yoshida et al. Clin. Immunol. Immunopathol. 82, 207-215 (1997), Thureau et al. Clin. Exp. Immunol. 109, 370-6 (1997), and Weiner et al. Res. Immunol. 148, 528-33 (1997). In particular certain routes of administration can cause tolerisation, such as oral, nasal or intraperitoneal. Particular products which cause tolerisation may be administered (e.g. in a composition which also comprises the agent) to the individual. Such products include cytokines, such as cytokines which favour a Th2 response (e.g. IL-4, TGF- β or IL-10). Products or agent may be administered at a dose which causes tolerisation.

Antagonists of T cells which recognise the agent can also be used to prevent or treat coeliac disease. Such an antagonist binds the TCR of the T cell (generally in the form of a complex with HLA-DQ2) but instead of causing normal functional activation causing an abnormal signal to be passed through the TCR intracellular

signalling cascade which causes the T cell to have decreased function activity (e.g. in response to recognition of an epitope, typically as measured by any suitable assay mentioned herein). Methods of causing antagonism are known in the art. In one embodiment the antagonist is a homologue of the epitopes mentioned above and may have any of the sequence, binding or other properties of the agent (particularly analogues). These antagonists typically differ from any of the above epitopes (which are capable of causing a normal antigen specific function in the T cell) by 1, 2, 3, 4 or more mutations (each of which may be a substitution, insertion or deletion). Such mutation are typically at the amino acid positions which contact the TCR.

Since the T cell immune response to the epitope of the invention in an individual is polyclonal more than one antagonist may need to be administered to cause antagonism of T cells of the response which have different TCRs. Therefore the antagonists may be administered in a composition which comprises at least 2, 4, 6 or more different antagonists, which each antagonise different T cells.

The invention also provides a method of identifying an antagonist of a T cell (which recognises the agent) comprising contacting a candidate substance with the T cell and detecting whether the substance causes a decrease in the ability of the T cell to undergo an antigen specific response (e.g. using any suitable assay mentioned herein), the detecting of any such decrease in said ability indicating that the substance is an antagonist.

In one embodiment the antagonists (including combinations of antagonists to a particular epitope) or tolerising (T cell and antibody tolerising) agents are present in a composition comprising at least 2, 4, 6 or more antagonists or agents which antagonise or tolerate to different epitopes of the invention, for example to the combinations of epitopes discussed above in relation to the agents which are a product comprising more than one substance.

Testing whether a composition is capable of causing coeliac disease

As mentioned above the invention provides a method of determining whether a composition is capable of causing coeliac disease comprising detecting the presence of a protein sequence which is capable of being modified by a

transglutaminase to as sequence comprising the agent or epitope of the invention (such transglutaminase activity may be a human intestinal transglutaminase activity). Typically this is performed by using a binding assay in which a moiety which binds to the sequence in a specific manner is contacted with the composition and the formation of sequence/moiety complex is detected and used to ascertain the presence of the agent. Such a moiety may be any suitable substance (or type of substance) mentioned herein, and is typically a specific antibody. Any suitable format of binding assay can be used (such as those mentioned herein).

In one embodiment the composition is contacted with at least 2, 5, 10 or more antibodies which are specific for epitopes of the invention from different gliadins, for example a panel of antibodies capable of recognising the combinations of epitopes discussed above in relation to agents of the invention which are a product comprising more than one substance.

The composition typically comprises material from a plant that expresses a gliadin which is capable of causing coeliac disease (for example any of the gliadins or plants mentioned herein). Such material may be a plant part, such as a harvested product (e.g. seed). The material may be processed products of the plant material (e.g. any such product mentioned herein), such as a flour or food that comprises the gliadin. The processing of food material and testing in suitable binding assays is routine, for example as mentioned in Kricka LJ, J. Biolumin. Chemilumin. 13, 189-93 (1998).

Binding assays

The determination of binding between any two substances mentioned herein may be done by measuring a characteristic of either or both substances that changes upon binding, such as a spectroscopic change.

The binding assay format may be a 'band shift' system. This involves determining whether the presence of one substance (such as a candidate substance) advances or retards the progress of the other substance during gel electrophoresis.

The format may be a competitive binding method which determines whether the one substance is able to inhibit the binding of the other substance to an agent

which is known to bind the other substance, such as a specific antibody.

Mutant gliadin proteins

5 The invention provides a gliadin protein in which an epitope sequence of the invention, or sequence which can be modified by a transglutaminase to provide such a sequence has been mutated so that it no longer causes, or is recognised by, a T cell response that recognises the epitope. In this context the term recognition refers to the TCR binding the epitope in such a way that normal (not antagonistic) antigen-specific functional activity of the T cell occurs.

10 Methods of identifying equivalent epitopes in other gliadins are discussed above. The wild type of the mutated gliadin is one which causes coeliac disease. Such a gliadin will have homology with SEQ ID NO:3, for example to the degree mentioned above (in relation to the analogue) across all of SEQ ID NO:3 or across 15, 30, 60, 100 or 200 contiguous amino acids of SEQ ID NO:3.

15 The mutated gliadin will not cause coeliac disease or will cause decreased symptoms of coeliac disease. Typically the mutation decreases the ability of the epitope to induce a T cell response. The mutated epitope may have a decreased binding to HLA-DQ2, a decreased ability to be presented by an APC or a decreased ability to bind to or to be recognised (i.e. cause antigen-specific functional activity) 20 by T cells that recognise the agent. The mutated gliadin or epitope will therefore show no or reduced recognition in any of the assays mentioned herein in relation to the diagnostic aspects of the invention.

The mutation may be one or more deletions, additions or substitutions of length 1 to 3, 4 to 6, 6 to 10, 11 to 15 or more in the epitope, for example across the 25 sequence SEQ ID NO:2 or its equivalent. Preferably the mutant gliadin has at least one mutation in the sequence SEQ ID NO:1. A preferred mutation is at position 65 in A-gliadin (or in an equivalent position in other gliadins). Typically the naturally occurring glutamine at this position is substituted to any of the amino acids shown in Table 3, preferably to histidine, tyrosine, tryptophan, lysine, proline, or arginine.

30 The invention thus also provides use of a mutation (such any of the mutations in any of the sequences discussed herein) in an epitope of a gliadin protein, which

epitope is an epitope of the invention, to decrease the ability of the gliadin protein to cause coeliac disease.

In one embodiment the mutated sequence is able to act as an antagonist. Thus the invention provides a protein that comprises a sequence which is able to bind
5 to a T cell receptor, which T cell receptor recognises an agent of the invention, and which sequence is able to cause antagonism of a T cell that carries such a T cell receptor.

The invention also provides proteins which are fragments of the above mutant gliadin proteins, which are at least 15 amino acids long (e.g. at least 30, 60, 100, 150,
10 200, or 250 amino acids long) and which comprise the mutations discussed above which decrease the ability of the gliadin to be recognised. Any of the mutant proteins (including fragments) mentioned herein may also be present in the form of fusion proteins, for example with other gliadins or with non-gliadin proteins.

The equivalent wild type protein to the mutated gliadin protein is typically
15 from a graminaceous monocotyledon, such as a plant of genus Triticum, e.g. wheat, rye, barley, oats or triticale. The protein is typically an α , $\alpha\beta$, β , γ or ω gliadin. The gliadin may be an A-gliadin.

Kits

20 The invention also provides a kit for carrying out the method comprising one or more agents and optionally a means to detect the recognition of the agent by the T cell. Typically the different agents are provided for simultaneous, separate or sequential use. Typically the means to detect recognition allows or aids detection based on the techniques discussed above.

25 Thus the means may allow detection of a substance secreted by the T cells after recognition. The kit may thus additionally include a specific binding moiety for the substance, such as an antibody. The moiety is typically specific for IFN- γ . The moiety is typically immobilised on a solid support. This means that after binding the moiety the substance will remain in the vicinity of the T cell which secreted it. Thus
30 'spots' of substance/moiety complex are formed on the support, each spot representing a T cell which is secreting the substance. Quantifying the spots, and

typically comparing against a control, allows determination of recognition of the agent.

5 The kit may also comprise a means to detect the substance/moiety complex. A detectable change may occur in the moiety itself after binding the substance, such as a colour change. Alternatively a second moiety directly or indirectly labelled for detection may be allowed to bind the substance/moiety complex to allow the determination of the spots. As discussed above the second moiety may be specific for the substance, but binds a different site on the substance than the first moiety.

10 The immobilised support may be a plate with wells, such as a microtitre plate. Each assay can therefore be carried out in a separate well in the plate.

The kit may additionally comprise medium for the T cells, detection moieties or washing buffers to be used in the detection steps. The kit may additionally comprise reagents suitable for the separation from the sample, such as the separation of PBMCs or T cells from the sample. The kit may be designed to allow detection of the T cells directly in the sample without requiring any separation of the components of the sample.

15 The kit may comprise an instrument which allows administration of the agent, such as intradermal or epidermal administration. Typically such an instrument comprises plaster, dressing or one or more needles. The instrument may allow ballistic delivery of the agent. The agent in the kit may be in the form of a pharmaceutical composition.

20 The kit may also comprise controls, such as positive or negative controls. The positive control may allow the detection system to be tested. Thus the positive control typically mimics recognition of the agent in any of the above methods. Typically in the kits designed to determine recognition *in vitro* the positive control is a cytokine. In the kit designed to detect *in vivo* recognition of the agent the positive control may be antigen to which most individuals should response.

25 The kit may also comprise a means to take a sample containing T cells from the host, such as a blood sample. The kit may comprise a means to separate mononuclear cells or T cells from a sample from the host.

Polynucleotides, cells, transgenic mammals and antibodies

The invention also provides a polynucleotide which is capable of expression to provide the agent or mutant gliadin proteins. Typically the polynucleotide is DNA or RNA, and is single or double stranded. The polynucleotide will preferably
5 comprise at least 50 bases or base pairs, for example 50 to 100, 100 to 500, 500 to 1000 or 1000 to 2000 or more bases or base pairs. The polynucleotide therefore comprises sequence which encodes the sequence of SEQ ID NO: 1 or 2 or any of the agents mentioned herein. To the 5' and 3' of this coding sequence the polynucleotide of the invention has sequence or codons which are different from the sequence or
10 codons 5' and 3' to these sequences in the corresponding gliadin gene.

5' and/or 3' to the sequence encoding the peptide the polynucleotide has coding or non-coding sequence. Sequence 5' and/or 3' to the coding sequence may comprise sequences which aid expression, such as transcription and/or translation, of the sequence encoding the agent. The polynucleotide may be capable of expressing
15 the agent prokaryotic or eukaryotic cell. In one embodiment the polynucleotide is capable of expressing the agent in a mammalian cell, such as a human, primate or rodent (e.g. mouse or rat) cell.

A polynucleotide of the invention may hybridise selectively to a polynucleotide that encodes SEQ ID NO:3 at a level significantly above background.
20 Selective hybridisation is typically achieved using conditions of medium to high stringency (for example 0.03M sodium chloride and 0.03M sodium citrate at from about 50°C to about 60°C). However, such hybridisation may be carried out under any suitable conditions known in the art (see Sambrook *et al* (1989), Molecular Cloning: A Laboratory Manual). For example, if high stringency is required, suitable
25 conditions include 0.2 x SSC at 60°C. If lower stringency is required, suitable conditions include 2 x SSC at 60°C.

Agents or proteins of the invention may be encoded by the polynucleotides described herein.

The polynucleotide may form or be incorporated into a replicable vector.
30 Such a vector is able to replicate in a suitable cell. The vector may be an expression vector. In such a vector the polynucleotide of the invention is operably linked to a

control sequence which is capable of providing for the expression of the polynucleotide. The vector may contain a selectable marker, such as the ampicillin resistance gene.

5 The polynucleotide or vector may be present in a cell. Such a cell may have been transformed by the polynucleotide or vector. The cell may express the agent. The cell will be chosen to be compatible with the said vector and may for example be a prokaryotic (bacterial), yeast, insect or mammalian cell. The polynucleotide or vector may be introduced into host cells using conventional techniques including calcium phosphate precipitation, DEAE-dextran transfection, or electroporation.

10 The invention provides processes for the production of the proteins of the invention by recombinant means. This may comprise (a) cultivating a transformed cell as defined above under conditions that allow the expression of the protein; and preferably (b) recovering the expressed polypeptide. Optionally, the polypeptide may be isolated and/or purified, by techniques known in the art.

15 The invention also provides TCRs which recognise (or bind) the agent, or fragments thereof which are capable of such recognition (or binding). These can be present in the any form mentioned herein (e.g. purity) discussed herein in relation to the protein of the invention. The invention also provides T cells which express such TCRs which can be present in any form (e.g. purity) discussed herein for the cells of the invention.

20 The invention also provides monoclonal or polyclonal antibodies which specifically recognise the agents (such as any of the epitopes of the invention) and which recognise the mutant gliadin proteins (and typically which do not recognise the equivalent wild-type gliadins) of the invention, and methods of making such antibodies. Antibodies of the invention bind specifically to these substances of the invention.

25 For the purposes of this invention, the term "antibody" includes antibody fragments such as Fv, F(ab) and F(ab)₂ fragments, as well as single-chain antibodies.

30 A method for producing a polyclonal antibody comprises immunising a suitable host animal, for example an experimental animal, with the immunogen and isolating immunoglobulins from the serum. The animal may therefore be inoculated

with the immunogen, blood subsequently removed from the animal and the IgG fraction purified. A method for producing a monoclonal antibody comprises immortalising cells which produce the desired antibody. Hybridoma cells may be produced by fusing spleen cells from an inoculated experimental animal with tumour cells (Kohler and Milstein (1975). *Nature* 256, 495-497).

An immortalized cell producing the desired antibody may be selected by a conventional procedure. The hybridomas may be grown in culture or injected intraperitoneally for formation of ascites fluid or into the blood stream of an allogenic host or immunocompromised host. Human antibody may be prepared by *in vitro* immunisation of human lymphocytes, followed by transformation of the lymphocytes with Epstein-Barr virus.

For the production of both monoclonal and polyclonal antibodies, the experimental animal is suitably a goat, rabbit, rat or mouse. If desired, the immunogen may be administered as a conjugate in which the immunogen is coupled, for example via a side chain of one of the amino acid residues, to a suitable carrier. The carrier molecule is typically a physiologically acceptable carrier. The antibody obtained may be isolated and, if desired, purified.

The polynucleotide, agent, protein or antibody of the invention, may carry a detectable label. Detectable labels which allow detection of the secreted substance by visual inspection, optionally with the aid of an optical magnifying means, are preferred. Such a system is typically based on an enzyme label which causes colour change in a substrate, for example alkaline phosphatase causing a colour change in a substrate. Such substrates are commercially available, e.g. from BioRad. Other suitable labels include other enzymes such as peroxidase, or protein labels, such as biotin, or radioisotopes, such as ^{32}P or ^{35}S . The above labels may be detected using known techniques.

Polynucleotides, agents, proteins, antibodies or cells of the invention may be in substantially purified form. They may be in substantially isolated form, in which case they will generally comprise at least 80% e.g. at least 90, 95, 97 or 99% of the polynucleotide, peptide, antibody, cells or dry mass in the preparation. The polynucleotide, agent, protein or antibody is typically substantially free of other

cellular components. The polynucleotide, agent, protein or antibody may be used in such a substantially isolated, purified or free form in the method or be present in such forms in the kit.

5 The invention also provides a transgenic mammal which expresses a TCR of the invention. This may be any of the mammals discussed herein (e.g. in relation to the production of the antibody). Preferably the mammal has, or is susceptible, to coeliac disease. The mammal may also express HLA-DQ2 and/or may be given a diet comprising a gliadin which cause coeliac disease (e.g. any of the gliadin proteins mentioned herein). Thus the mammal may act as an animal model for coeliac
10 disease.

The invention also provides a method of identifying a product which is therapeutic for coeliac disease comprising administering a candidate substance to a mammal of the invention which has, or which is susceptible to, coeliac disease and determining whether substance prevents or treats coeliac disease in the mammal, the
15 prevention or treatment of coeliac disease indicating that the substance is a therapeutic product. Such a product may be used to treat or prevent coeliac disease.

The invention provides therapeutic (including prophylactic) agents or diagnostic substances (the agents, proteins and polynucleotides of the invention). These substances are formulated for clinical administration by mixing them with a
20 pharmaceutically acceptable carrier or diluent. For example they can be formulated for topical, parenteral, intravenous, intramuscular, subcutaneous, intraocular, intradermal, epidermal or transdermal administration. The substances may be mixed with any vehicle which is pharmaceutically acceptable and appropriate for the desired route of administration. The pharmaceutically carrier or diluent for injection
25 may be, for example, a sterile or isotonic solution such as Water for Injection or physiological saline, or a carrier particle for ballistic delivery.

The dose of the substances may be adjusted according to various parameters, especially according to the agent used; the age, weight and condition of the patient to be treated; the mode of administration used; the severity of the condition to be
30 treated; and the required clinical regimen. As a guide, the amount of substance administered by injection is suitably from 0.01 mg/kg to 30 mg/kg, preferably from

0.1 mg/kg to 10 mg/kg.

The routes of administration and dosages described are intended only as a guide since a skilled practitioner will be able to determine readily the optimum route of administration and dosage for any particular patient and condition.

5 The substances of the invention may thus be used in a method of treatment of the human or animal body, or in a diagnostic method practised on the human body. In particular they may be used in a method of treating or preventing coeliac disease. The invention also provide the agents for use in a method of manufacture of a medicament for treating or preventing coeliac disease. Thus the invention provides a
10 method of preventing or treating coeliac disease comprising administering to a human in need thereof a substance of the invention (typically a non-toxic effective amount thereof).

 The agent of the invention can be made using standard synthetic chemistry techniques, such as by use of an automated synthesizer. The agent may be made
15 from a longer polypeptide e.g. a fusion protein, which polypeptide typically comprises the sequence of the peptide. The peptide may be derived from the polypeptide by for example hydrolysing the polypeptide, such as using a protease; or by physically breaking the polypeptide. The polynucleotide of the invention can be made using standard techniques, such as by using a synthesiser.

20 Plant cells and plants that express mutant gliadin proteins

 The cell of the invention may be a plant cell, such as a cell of a graminaceous monocotyledonous species. The species may be one whose wild-type form expresses gliadins, such as any of the gliadin proteins mentioned herein (including gliadins
25 ~~with any degree of~~ homology to SEQ ID NO:3 mentioned herein). Such a gliadin may cause coeliac disease in humans. The cell may be of wheat, maize, oats, rye, rice, barley, triticale, sorghum, or sugar cane. Typically the cell is of the Triticum genus, such as aestivum, spelta, polonicum or monococcum.

 The plant cell of the invention is typically one which does not express a wild-
30 type gliadin (such as any of the gliadins mentioned herein which may cause coeliac disease), or one which does not express a gliadin comprising a sequence that can be

recognised by a T cell that recognises the agent. Thus if the wild-type plant cell did express such a gliadin then it may be engineered to prevent or reduce the expression of such a gliadin or to change the amino acid sequence of the gliadin so that it no longer causes coeliac disease (typically by no longer expressing the epitope of the invention).

This can be done for example by introducing mutations into 1, 2, 3 or more or all of such gliadin genes in the cell, for example into coding or non-coding (e.g. promoter regions). Such mutations can be any of the type or length of mutations discussed herein (e.g in relation to homologous proteins). The mutations can be introduced in a directed manner (e.g using site directed mutagenesis or homologous recombination techniques) or in a random manner (e.g. using a mutagen, and then typically selecting for mutagenised cells which no longer express the gliadin (or a gliadin sequence which causes coeliac disease)).

The polynucleotide which is present in (or which was transformed into) the plant cell will generally comprise promoter capable of expressing the mutant gliadin protein the plant cell. Depending on the pattern of expression desired, the promoter may be constitutive, tissue- or stage-specific; and/or inducible. For example, strong constitutive expression in plants can be obtained with the CAMV 35S, Rubisco ssu, or histone promoters. Also, tissue-specific or stage-specific promoters may be used to target expression of protein of the invention to particular tissues in a transgenic plant or to particular stages in its development. Thus, for example seed-specific, root-specific, leaf-specific, flower-specific etc promoters may be used. Seed-specific promoters include those described by Dalta *et al* (Biotechnology Ann. Rev. (1997), 3, pp.269-296). Particular examples of seed-specific promoters are napin promoters (EP-A-0 255, 378), phaseolin promoters, glutenine promoters, helianthene promoters (WO92/17580), albumin promoters (WO98/45460), oleosin promoters (WO98/45461) and ATS1 and ATS3 promoters (PCT/US98/06798).

The cell may be in any form. For example, it may be an isolated cell, e.g. a protoplast, or it may be part of a plant tissue, e.g. a callus, or a tissue excised from a plant, or it may be part of a whole plant. The cell may be of any type (e.g of any type of plant part). For example, an undifferentiated cell, such as a callus cell; or a

differentiated cell, such as a cell of a type found in embryos, pollen, roots, shoots or leaves. Plant parts include roots; shoots; leaves; and parts involved in reproduction, such as pollen, ova, stamens, anthers, petals, sepals and other flower parts.

The invention provides a method of obtaining a transgenic plant cell
5 comprising transforming a plant cell with a polynucleotide or vector of the invention to give a transgenic plant cell. Any suitable transformation method may be used (in the case of wheat the techniques disclosed in Vasil V et al, Biotechnology 10, 667-674 (1992) may be used). Preferred transformation techniques include electroporation of plant protoplasts and particle bombardment. Transformation may
10 thus give rise to a chimeric tissue or plant in which some cells are transgenic and some are not.

The cell of the invention or thus obtained cell may be regenerated into a transgenic plant by techniques known in the art. These may involve the use of plant growth substances such as auxins, giberellins and/or cytokinins to stimulate the
15 growth and/or division of the transgenic cell. Similarly, techniques such as somatic embryogenesis and meristem culture may be used. Regeneration techniques are well known in the art and examples can be found in, e.g. US 4,459,355, US 4,536,475, US 5,464,763, US 5, 177,010, US 5, 187,073, EP 267,159, EP 604, 662, EP 672, 752, US 4,945,050, US 5,036,006, US 5,100,792, US 5,371,014, US 5,478,744, US
20 5,179,022, US 5,565,346, US 5,484,956, US 5,508,468, US 5,538,877, US 5,554,798, US 5,489,520, US 5,510,318, US 5,204,253, US 5,405,765, EP 442,174, EP 486,233, EP 486,234, EP 539,563, EP 674,725, WO91/02071 and WO 95/06128.

In many such techniques, one step is the formation of a callus, i.e. a plant tissue comprising expanding and/or dividing cells. Such calli are a further aspect of
25 the invention as are other types of plant cell cultures and plant parts. Thus, for example, the invention provides transgenic plant tissues and parts, including embryos, meristems, seeds, shoots, roots, stems, leaves and flower parts. These may be chimeric in the sense that some of their cells are cells of the invention and some are not. Transgenic plant parts and tissues, plants and seeds of the invention may be
30 of any of the plant species mentioned herein.

Regeneration procedures will typically involve the selection of transformed

cells by means of marker genes.

The regeneration step gives rise to a first generation transgenic plant. The invention also provides methods of obtaining transgenic plants of further generations from this first generation plant. These are known as progeny transgenic plants.

5 Progeny plants of second, third, fourth, fifth, sixth and further generations may be obtained from the first generation transgenic plant by any means known in the art.

Thus, the invention provides a method of obtaining a transgenic progeny plant comprising obtaining a second-generation transgenic progeny plant from a first-generation transgenic plant of the invention, and optionally obtaining transgenic
10 plants of one or more further generations from the second-generation progeny plant thus obtained.

Progeny plants may be produced from their predecessors of earlier generations by any known technique. In particular, progeny plants may be produced by:

15 obtaining a transgenic seed from a transgenic plant of the invention belonging to a previous generation, then obtaining a transgenic progeny plant of the invention belonging to a new generation by growing up the transgenic seed; and/or

propagating clonally a transgenic plant of the invention belonging to a previous
20 generation to give a transgenic progeny plant of the invention belonging to a new generation; and/or

crossing a first-generation transgenic plant of the invention belonging to a previous generation with another compatible plant to give a transgenic progeny plant of the
25 invention belonging to a new generation; and optionally

obtaining transgenic progeny plants of one or more further generations from the progeny plant thus obtained.

30 These techniques may be used in any combination. For example, clonal propagation and sexual propagation may be used at different points in a process that

gives rise to a transgenic plant suitable for cultivation. In particular, repetitive back-crossing with a plant taxon with agronomically desirable characteristics may be undertaken. Further steps of removing cells from a plant and regenerating new plants therefrom may also be carried out.

5 Also, further desirable characteristics may be introduced by transforming the cells, plant tissues, plants or seeds, at any suitable stage in the above process, to introduce desirable coding sequences other than the polynucleotides of the invention. This may be carried out by the techniques described herein for the introduction of polynucleotides of the invention.

10 For example, further transgenes may be selected from those coding for other herbicide resistance traits, e.g. tolerance to: Glyphosate (e.g. using an EPSP synthase gene (e.g. EP-A-0 293,358) or a glyphosate oxidoreductase (WO 92/000377) gene); or tolerance to fosametin; a dihalobenzonitrile; glufosinate, e.g. using a phosphinothrycin acetyl transferase (PAT) or glutamine synthase gene
15 (cf. EP-A-0 242,236); asulam, e.g. using a dihydropteroate synthase gene (EP-A-0 369,367); or a sulphonylurea, e.g. using an ALS gene); diphenyl ethers such as acifluorfen or oxyfluorfen, e.g. using a protoporphyrinogen oxidase gene); an oxadiazole such as oxadiazon; a cyclic imide such as chlorophthalim; a phenyl pyrazole such as TNP, or a phenopylate or carbamate analogue thereof.

20 Similarly, genes for beneficial properties other than herbicide tolerance may be introduced. For example, genes for insect resistance may be introduced, notably genes encoding *Bacillus thuringiensis* (Bt) toxins. Likewise, genes for disease resistance may be introduced, e.g. as in WO91/02701 or WO95/06128.

25 Typically, a protein of the invention is expressed in a plant of the invention. Depending on the promoter used, this expression may be constitutive or inducible. Similarly, it may be tissue- or stage-specific, i.e. directed towards a particular plant tissue (such as any of the tissues mentioned herein) or stage in plant development.

30 The invention also provides methods of obtaining crop products by harvesting, and optionally processing further, transgenic plants of the invention. By crop product is meant any useful product obtainable from a crop plant.

Products that contain mutant gliadin proteins

The invention provides a product that comprises the mutant gliadin proteins. This is typically derived from or comprise plant parts from plants mentioned herein which express the mutant gliadin proteins. Such a product may be obtainable
5 directly by harvesting or indirectly, by harvesting and further processing the plant of the invention. Directly obtainable products include grains. Alternatively, such a product may be obtainable indirectly, by harvesting and further processing. Examples of products obtainable by further processing are flour or distilled alcoholic beverages; food products made from directly obtained or further processed material,
10 e.g. baked products (e.g. bread) made from flour. Typically such food products, which are ingestible and digestible (i.e. non-toxic and of nutrient value) by human individuals.

The invention is illustrated by the following Examples:

Example 1

15 We carried out epitope mapping in Coeliac disease by using a set of 51 synthetic 15-mer peptides that span the complete sequence of a fully characterized a-gliadin, "A-gliadin" (see Table 1). A-Gliadin peptides were also individually treated with tTG to generate products that might mimic those produced in vivo³. We also sought to study Coeliac disease patients at the point of initiation of disease relapse to
20 avoid the possibility that epitope "spreading" or "exhaustion" may have occurred, as described in experimental infectious and autoimmune diseases.

Clinical and A-gliadin specific T cell responses with 3 and 10 day bread challenge

25 In a pilot study, two subjects with Coeliac disease in remission, defined by absence of serum anti-endomysial antibody (EMA), on a gluten free diet were fed four slices of standard gluten-containing white bread daily in addition to their usual gluten free diet. Subject 1 ceased bread because of abdominal pain, mouth ulcers and mild diarrhoea after three days, but Subject 2 continued for 10 days with only mild nausea at one week. The EMA became positive in Subject 2 one week after the
30 bread challenge, indicating the bread used had caused a relapse of Coeliac disease. But in Subject 1, EMA remained negative up to two months after bread challenge. In

both subjects, symptoms that appeared with bread challenge resolved within two days after returning to gluten free diet.

PBMC responses in IFN γ ELISPOT assays to A-gliadin peptides were not found before or during bread challenge. But from the day after bread withdrawal (Day 4) in Subject 1 a single pool of 5 overlapping peptides spanning A-gliadin 51-85 (Pool 3) treated with tTG showed potent IFN γ responses (see Figure 1a). In Subject 1, the PBMC IFN γ response to A-gliadin peptide remained targeted to Pool 3 alone and was maximal on Day 8. The dynamics and magnitude of the response to Pool 3 was similar to that elicited by α -chymotrypsin digested gliadin. PBMC IFN γ responses to tTG-treated Pool 3 were consistently 5 to 12-fold greater than Pool 3 not treated with tTG, and responses to α -chymotrypsin digested gliadin were 3 to 10-fold greater if treated with tTG. In Subject 2, Pool 3 treated with tTG was also the only immunogenic set of A-gliadin peptides on Day 8, but this response was weaker than Subject 1, was not seen on Day 4 and by Day 11 the response to Pool 3 had diminished and other tTG-treated pools of A-gliadin peptides elicited stronger IFN α responses (see Figure 1b).

The pilot study indicated that the initial T cell response in these Coeliac disease subjects was against a single tTG-treated A-gliadin pool of five peptides and was readily measured in peripheral blood. But if antigen exposure is continued for ten days instead of three, T cell responses to other A-gliadin peptides appear, consistent with epitope spreading.

Coeliac disease-specific IFN-g induction by tTG-treated A-gliadin peptides

In five out of six further Coeliac disease subjects on gluten free diet (see Table 1), bread challenge for three days identified tTG-treated peptides in Pool 3, and in particular, peptides corresponding to 56-70 (12) and 60-75 (13) as the sole A-gliadin components eliciting IFN γ from PBMC (see Figure 2). IL-10 ELISPOT assays run in parallel to IFN γ ELISPOT showed no IL-10 response to tTG-treated peptides 12 or 13. In one subject, there were no IFN γ responses to any A-gliadin peptide or α -chymotrypsin digested gliadin before, during or up to four days after bread challenge. In none of these Coeliac disease subjects did EMA status change

from baseline when measured for up to two months after bread challenge.

PBMC from four healthy, EMA-negative subjects with the HLA-DQ alleles $\alpha 1^*0501$, $\beta 1^*0201$ (ages 28-52, 2 females) who had been challenged for three days with bread after following a gluten free diet for one month, showed no IFN γ responses above the negative control to any of the A-gliadin peptides with or without tTG treatment. Thus, induction of IFN γ in PBMC to tTG-treated Pool 3 and A-gliadin peptides 56-70 (12) and 60-75 (13) were Coeliac disease specific (7/8 vs 0/4, $p < 0.01$ by Chi-squared analysis).

Fine mapping of the minimal A-gliadin T cell epitope

tTG-treated peptides representing truncations of A-gliadin 56-75 revealed that the same core peptide sequence (QPQLP) was essential for antigenicity in all of the five Coeliac disease subjects assessed (see Figure 3). PBMC IFN γ responses to tTG-treated peptides spanning this core sequence beginning with the 7-mer PQPQLPY and increasing in length, indicated that the tTG-treated 17-mer QLQPFPPQPQLPYQPQS (A-gliadin 57-73) possessed optimal activity in the IFN γ ELISPOT (see Figure 4).

Deamidation of Q65 by tTG generates the immunodominant T cell epitope in A-gliadin

HPLC analysis demonstrated that tTG treatment of A-gliadin 56-75 generated a single product that eluted marginally later than the parent peptide. Amino acid sequencing indicated that out of the six glutamine (Q) residues contained in A-gliadin 56-75, Q65 was preferentially deamidated by tTG (see Figure 5). Bioactivity of peptides corresponding to serial expansions from the core A-gliadin 62-68 sequence in which glutamate (E) replaced Q65, was equivalent to the same peptides with Q65 after tTG-treatment (see Figure 4a). Replacement of Q57 and Q72 by E together or alone, with E65 did not enhance antigenicity of the 17-mer in the three Coeliac disease subjects studied (see Figure 6). Q57 and Q72 were investigated because glutamine residues followed by proline in gliadin peptides are not deamidated by tTG in vitro (W. Vader et al, Proceedings 8th International

Symposium Coeliac Disease). Therefore, the immunodominant T cell epitope was defined as QLQPF~~P~~QPELPYPQPQS.

Immunodominant T cell epitope response is DQ2-restricted and CD4 dependent

5 In two Coeliac disease subjects homozygous for HLA-DQ $\alpha 1^*0501$, $\beta 1^*0201$, anti-DQ monoclonal antibody blocked the ELISPOT IFN γ response to tTG-treated A-gliadin 56-75, but anti-DP and -DR antibody did not (see Figure 7). Anti-CD4 and anti-CD8 magnetic bead depletion of PBMC from two Coeliac disease subjects indicated the IFN γ response to tTG-treated A-gliadin 56-75 is CD4 T cell-mediated.
10

Discussion

 In this study we describe a rather simple dietary antigen challenge using standard white bread to elicit a transient population of CD4 T cells in peripheral
15 blood of Coeliac disease subjects responsive to a tTG-treated A-gliadin 17-mer with the sequence: QLQPF~~P~~QPELPYPQPQS (residues 57-73). The immune response to A-gliadin 56-75 (Q-E65) is restricted to the Coeliac disease-associated HLA allele, DQ $\alpha 1^*0501$, $\beta 1^*0201$. Tissue transglutaminase action in vitro selectively deamidates Q65. Elicited peripheral blood IFN γ responses to synthetic A-gliadin
20 peptides with the substitution Q-E65 is equivalent to tTG-treated Q65 A-gliadin peptides; both stimulate up to 10-fold more T cells in the IFN γ ELISPOT than unmodified Q65 A-gliadin peptides.

 We have deliberately defined this Coeliac disease-specific T cell epitope using in vivo antigen challenge and short-term ex vivo immune assays to avoid the
25 possibility of methodological artifacts that may occur with the use of T cell clones in epitope mapping. Our findings indicate that peripheral blood T cell responses to ingestion of gluten are rapid but short-lived and can be utilized for epitope mapping. In vivo antigen challenge has also shown there is a temporal hierarchy of immune responses to A-gliadin peptides; A-gliadin 57-73 modified by tTG not only elicits the
30 strongest IFN γ response in PBMC but it is also the first IFN γ response to appear.

 Because we have assessed only peptides spanning A-gliadin, there may be

other epitopes in other gliadins of equal or greater importance in the pathogenesis of Coeliac disease. Indeed, the peptide sequence at the core of the epitope in A-gliadin that we have identified (PQPQLPY) is shared by several other gliadins (SwissProt and Trembl accession numbers: P02863, Q41528, Q41531, Q41533, Q9ZP09, P04722, P04724, P18573). However, A-gliadin peptides that have previously been shown to possess bioactivity in biopsy challenge and in vivo studies (for example: 31-43, 44-55, and 206-217)^{4,5} did not elicit IFN γ responses in PBMC following three day bread challenge in Coeliac disease subjects. These peptides may be "secondary" T cell epitopes that arise with spreading of the immune response.

Example 2

The effect on T cell recognition of substitutions in the immunodominant epitope

The effect of substituting the glutamate at position 65 in the 57-73 A-gliadin epitope was determined by measuring peripheral blood responses against the substituted epitopes in an IFN γ ELISPOT assay using synthetic peptides (at 50 μ g/ml). The responses were measured in 3 Coeliac disease subjects 6 days after commencing gluten challenge (4 slices bread daily for 3 days). Results are shown in table 3 and Figure 8. As can be seen substitution of the glutamate to histidine, tyrosine, tryptophan, lysine, proline or arginine stimulated a response whose magnitude was less than 10% of the magnitude of the response to the immunodominant epitope. Thus mutation of A-gliadin at this position could be used to produce a mutant gliadin with reduce or absent immunoreactivity.

Example 3

Testing the immunoreactivity of equivalent peptides from other naturally occurring gliadins

The immunoreactivity of equivalent peptides from other naturally occurring wheat gliadins was assessed using synthetic peptides corresponding to the naturally occurring sequences which were then treated with transglutaminase. These peptides were tested in an ELISPOT in the same manner and with PBMCs from the same

subjects as described in Example 2. At least five of the peptides show immunoreactivity comparable to the A-gliadin 57-73 E65 peptide (after transglutaminase treatment) indicating that other gliadin proteins in wheat are also likely to induce this Coeliac disease-specific immune response (Table 4 and Figure 9).

Methods

Subjects: Patients used in the study attended a Coeliac Clinic in Oxford, United Kingdom. Coeliac disease was diagnosed on the basis of typical small intestinal histology, and normalization of symptoms and small intestinal histology with gluten free diet.

Tissue typing: Tissue typing was performed using DNA extracted from EDTA-anticoagulated peripheral blood. HLA-DQA and DQB genotyping was performed by PCR using sequence-specific primer mixes⁶⁻⁸.

Anti-endomysial antibody assay: EMA were detected by indirect immunofluorescence using patient serum diluted 1:5 with monkey oesophagus, followed by FITC-conjugated goat anti-human IgA. IgA was quantitated prior to EMA, none of the subjects were IgA deficient.

Antigen Challenge: Coeliac disease subjects following a gluten free diet, consumed 4 slices of gluten-containing bread (50g/slice, Sainsbury's "standard white sandwich bread") daily for 3 or 10 days. EMA was assessed the week before and up to two months after commencing the bread challenge. Healthy subjects who had followed a gluten free diet for four weeks, consumed their usual diet including four slices of gluten-containing bread for three days, then returned to gluten free diet for a further six days.

IFN γ and IL-10 ELISPOT: PBMC were prepared from 50-100 ml of venous blood by Ficoll-Hypaque density centrifugation. After three washes, PBMC were

resuspended in complete RPMI containing 10% heat inactivated human AB serum. ELISPOT assays for single cell secretion of IFN γ and IL-10 were performed using commercial kits (Mabtech; Stockholm, Sweden) with 96-well plates (MAIP-S-45; Millipore, Bedford, MA) according to the manufacturers instructions (as described elsewhere⁹) with 2.5×10^5 (IFN γ) or $0.4-1 \times 10^5$ (IL-10) PBMC in each well. Peptides were assessed in duplicate wells, and Mycobacterium tuberculosis purified protein derivative (PPD RT49) (Serum Institute; Copenhagen, Denmark) (20 μ g/ml) was included as a positive control in all assays.

Peptides: Synthetic peptides were purchased from Research Genetics (Huntsville, Alabama) Mass-spectroscopy and HPLC verified peptides' authenticity and >70% purity. Digestion of gliadin (Sigma; G-3375) (100 mg/ml) with α -chymotrypsin (Sigma; C-3142) 200:1 (w/w) was performed at room temperature in 0.1 M NH₄HCO₃ with 2M urea and was halted after 24 h by heating to 98°C for 10 minutes. After centrifugation (13 000g, 10 minutes), the gliadin digest supernatant was filter-sterilized (0.2 μ m). Digestion of gliadin was verified by SDS-PAGE and protein concentration assessed. α -Chymotrypsin-digested gliadin (640 μ g/ml) and synthetic gliadin peptides (15-mers: 160 μ g/ml, other peptides: 0.1 mM) were individually treated with tTG (Sigma; T-5398) (50 μ g/ml) in PBS + CaCl₂ 1 mM for 2 h at 37°C. Peptides and peptide pools were aliquotted into sterile 96-well plates and stored frozen at -20°C until use.

Amino acid sequencing of peptides: Reverse phase HPLC was used to purify the peptide resulting from tTG treatment of A-gliadin 56-75. A single product was identified and subjected to amino acid sequencing (automated sequencer Model 494A, Applied Biosystems, Foster City, California). The sequence of unmodified G56-75 was confirmed as: LQLQFPQPQLPYQPQSFP, and tTG treated G56-75 was identified as: LQLQFPQPELPYQPQSFP. Deamidation of glutamyl residues was defined as the amount (pmol) of glutamate recovered expressed as a percent of the combined amount of glutamine and glutamate recovered in cycles 2, 4, 8, 10, 15 and 17 of the amino acid sequencing. Deamidation attributable to tTG was defined

as (% deamidation of glutamine in the tTG treated peptide - % deamidation in the untreated peptide) / (100 - % deamidation in the untreated peptide).

CD4/CD8 and HLA Class II Restriction: Anti-CD4 or anti-CD8 coated magnetic
5 beads (Dyna, Oslo, Norway) were washed four times with RPMI then incubated
with PBMC in complete RPMI containing 10% heat inactivated human AB serum
(5×10^6 cells/ml) for 30 minutes on ice. Beads were removed using a magnet and
cells remaining counted. In vivo HLA-class II restriction of the immune response to
tTG-treated A-gliadin 56-75 was established by incubating PBMC (5×10^6 cells/ml)
10 with anti-HLA-DR (L243), -DQ (L2), and -DP (B7.21) monoclonal antibodies (10
 μ g/ml) at room temperature for one hour prior to the addition of peptide.

References

- 15 1. Molberg O, et al. Nature Med. 4, 713-717 (1998).
2. Quarsten H, et al. Eur. J. Immunol. 29, 2506-2514 (1999).
3. Greenberg CS et al. FASEB 5, 3071-3077 (1991).
4. Mantzaris G, Jewell D. Scand. J. Gastroenterol. 26, 392-398 (1991).
5. Mauri L, et al. Scand. J. Gastroenterol. 31, 247-253 (1996).
- 20 6. Bunce M, et al. Tissue Antigens 46, 355-367 (1995).
7. Olerup O, et al. Tissue antigens 41, 119-134 (1993).
8. Mullighan CG, et al. Tissue-Antigens. 50, 688-92 (1997).
9. Plebanski M et al. Eur. J. Immunol. 28, 4345-4355 (1998).

Table 1. A-Gliadin protein sequence (based on amino acid sequencing)

VRVPVQLQP QNPSQQQPQE QVPLVQQQQF PGQQQQFPPQ QPYPQPQFP SQQPYLQLQP FPQPQLPYPQ
1 11 21 31 41 51 61
PQSFPQQPY PQPQOYSQP QQPISQQQAQ QQQQQQQQQQ QQQILQQILQ QQLPCMDVV LQQHNIAHAR
71 81 91 101 111 121 131
SQVLQQSTYQ LLQELCCQHL WQIPEQSQCQ AIHNVVHAII LHQQKQKQQQ PSSQVSFQQP LQQYP LGQGS
141 151 161 171 181 191 201
FRPSQQNPQA QGSVQPQQLP QFEEIRNLAL QTLPAMCNVY IAPYCTIAPF GIFGTN
211 221 231 241 251 261

Table 2. Coeliac disease subjects studied

	Age Sex	Gluten free diet	HLA-DQ2	Bread challenge	Symptoms with bread
1	64 f	14 yr	Homozygote	3 days	Abdominal pain, lethargy, mouth ulcers, diarrhoea
2	57 m	1 yr	Heterozygote	10 days	Lethargy, nausea
3	35 f	7 yr	Heterozygote	3 days	Nausea
4	36 m	6 wk	Homozygote	3 days	Abdominal pain, mouth ulcers, diarrhoea
5	26 m	19 yr	Heterozygote	3 days	None
6	58 m	35 yr	Heterozygote	3 days	None
7	55 m	1 yr	Heterozygote	3 days	Diarrhoea
8	48 f	15 yr	Homozygote	3 days	Abdominal pain, diarrhoea

Table 3

Aminoacid at position 65	Range	Mean
Glutamate	(100)	100%
Asparagine	(50-84)	70%
Aspartate	(50-94)	65%
Alanine	(44-76)	64%
Cysteine	(45-83)	62%
Serine	(45-75)	62%
Valine	(24-79)	56%
Threonine	(46-66)	55%
Glycine	(34-47)	40%
Leucine	(8-46)	33%
Glutamine	(16-21)	19%
Isoleucine	(3-25)	14%
Methionine	(3-32)	14%
Phenylalanine	(0-33)	12%
Histidine	(0-13)	8%
Tyrosine	(0-17)	8%
Tryptophan	(0-17)	8%
Lysine	(0-11)	4%
Proline	(0-4)	2%
Arginine	(0-2)	1%

Table 4

Elisopt response No TG TG	Peptide sequence	Corresponding residues in gliadin protein sequences (Accession no.)	
8 (1-13)	QLQPFPPQLPYPPQPS	57-73	α -Gliadin (T. aestivum) Q41545
	QLQPFPPQLPYPPQPS	57-73	α -Gliadin (T. aestivum) Q41545
5 (1-7)	QLQPFPPQLPYSQPQP	77-93	α/β -Gliadin precursor (Triticum. aestivum) P02863
		76-92	α -Gliadin (T. aestivum) Q41528
		77-93	α -Gliadin storage protein (T. aestivum) Q41531
		57-73	α -Gliadin mature peptide (T. aestivum) Q41533
		77-93	α -Gliadin precursor (T. spelta) Q9ZP09
12 (0-20)	QLQPFPPQLPYPPQPF	77-93	α/β -Gliadin A-II precursor (T. aestivum) P0472
19 (0-33)	QLQPFPPQLPYPPQQL	77-93	α/β -Gliadin A-IV precursor (T. aestivum) P04724
		77-93	α/β -Gliadin MM1 precursor (T. aestivum) P18573
3 (0-7)	PQLPYPPQLPYPPQPF	84-100	α/β -Gliadin A-IV precursor (T. aestivum) P04724
ND	PQLPYPPQLPYPPQQL	84-100	α/β -Gliadin MM1 precursor (T. aestivum) P18573
0 (0-1)	QLQPFPPQLPYSQPQP	77-93	α/β -Gliadin A-I precursor (T. aestivum) P04721
		77-93	α -Gliadin (T. aestivum) Q41509
0 (0-0)	QLQPFPPQLPYSQPQP	77-93	α -Gliadin storage protein (T. aestivum) Q41530
ND	PQPQFPFPQLPYPPQTP	77-93	α/β -Gliadin A-III precursor (T. aestivum) P04723
17 (0-40)	PQPQFPFPQLPYPPQPS	82-98	α/β -Gliadin A-V precursor (T. aestivum) P04725
10 (0-30)	PQPQFPFPQLPYPPPP	82-98	α/β -Gliadin clone PW1215 precursor (T. aestivum) P04726
		82-98	α/β -Gliadin (T. urartu) Q41632
10 (0-30)	PQPQFPLQLPYPPQPS	79-95	α/β -Gliadin clone PW8142 precursor (T. aestivum) P04726
		79-95	α -Gliadin (T. aestivum) Q41529
		79-95	α/β -Gliadin precursor (T. aestivum) Q41546

CLAIMS

1. A method of diagnosing coeliac disease, or susceptibility to coeliac disease, in an individual comprising:
 - (a) contacting a sample from the host with an agent selected from
 - (i) the epitope comprising sequence which is: SEQ ID NO:1 or 2, or an equivalent sequence from a naturally occurring homologue of the gliadin represented by SEQ ID NO:3,
 - (ii) an epitope comprising sequence comprising: SEQ ID NO:1, or an equivalent sequence from a naturally occurring homologue of the gliadin represented by SEQ ID NO:3, which epitope is an isolated oligopeptide derived from a gliadin protein,
 - (iii) an analogue of (i) or (ii) which is capable of being recognised by a T cell receptor that recognises (i) or (ii), which in the case of a peptide analogue is not more than 50 amino acids in length, or
 - (iv) a product comprising two or more agents as defined in (i), (ii) or (iii), and
 - (b) determining *in vitro* whether T cells in the sample recognise the agent; recognition by the T cells indicating that the individual has, or is susceptible to, coeliac disease.
2. Use of an agent as defined in claim 1 for the preparation of a diagnostic means for use in a method of diagnosing coeliac disease, or susceptibility to coeliac disease, in an individual, said method comprising determining whether T cells of the individual recognise the agent, recognition by the T cells indicating that the individual has, or is susceptible to, coeliac disease.
3. A method or use according to claim 1 or 2 wherein the agent is an analogue (iii) which comprises (i) or (ii) bound to (a) an HLA molecule, or (b) a fragment of an HLA molecule capable of binding (i) or (ii).
4. A method or use according to claim 3 wherein the HLA molecule or fragment

is in a complex comprising four HLA molecules or fragments of HLA molecules.

5. Use according to claim 2, 3 or 4 wherein the method comprises administering the agent to the skin of an individual and detecting the presence of inflammation at the site of administration, the detection of inflammation indicating that the T cells of the individual recognise the agent.
6. A method according to claim 1, 3 or 4 wherein the sample is blood sample.
7. A method according to claim 1, 3, 4 or 6 wherein the T cells are not restimulated in antigen specific manner *in vitro* before the said determining.
8. A method or use according to any one of the preceding claims in which the recognition of the agent by the T cells is determined by detecting the secretion of a cytokine from the T cells.
9. A method or use according to claim 8 in which the cytokine is IFN- γ .
10. A method or use according to claim 8 or claim 9 in which the cytokine is detected by allowing the cytokine to bind to an immobilised antibody specific to the cytokine and then detecting the presence of the antibody/cytokine complex.
11. A method or use according to any one of claims 1 to 7 wherein said determining is done by measuring whether the agent binds the T cell receptor.
12. A method for identifying an analogue as defined in a claim 1, 3 or 4 comprising determining whether a candidate substance is recognised by a T cell receptor that recognises an epitope comprising sequence as defined in claim 1, recognition of the substance indicating that the substance is an

analogue.

13. A method of diagnosing coeliac disease, or susceptibility to coeliac disease, in an individual comprising determining the presence of an antibody that binds to an epitope of an epitope comprising sequence as defined in claim 1 in a sample from the individual, the presence of the antibody indicating that the individual has, or is susceptible to, coeliac disease.
14. An agent as defined in claim 1, optionally in association with a carrier, for use in a method of treating or preventing coeliac disease by tolerising T cells which recognise the agent.
15. An antagonist of a T cell which has a T cell receptor as defined in claim 1(iii), optionally in association with a carrier, for use in a method of treating or preventing coeliac disease by antagonising such T cells.
16. An agent as defined in claim 1 or an analogue that binds an antibody as defined in claim 13 for use in a method of treating or preventing coeliac disease in an individual by tolerising the individual to prevent the production of such an antibody.
17. A method of determining whether a composition is capable of causing coeliac disease comprising determining whether a protein capable of being modified by a transglutaminase to an oligopeptide sequence as defined in claim 1 is present in the composition, the presence of the protein indicating that the composition is capable of causing coeliac disease.
18. A method according to claim 17 wherein the said determining is done by contacting the composition with an antibody specific for the sequence which is capable of being modified to the oligopeptide sequence, binding of the antibody to a protein in the composition indicating the composition is capable

of causing coeliac disease.

19. A mutant gliadin protein whose wild-type sequence can be modified by a transglutaminase to a sequence that comprises an epitope comprising sequence as defined in claim 1, but which mutant gliadin protein has been modified in such a way that it does not contain sequence which can be modified by a transglutaminase to a sequence that comprises an epitope comprising sequence as defined in claim 1; or a fragment of such a mutant gliadin protein which is at least 15 amino acids long and which comprises sequence which has been modified in said way.
20. A protein that comprises a sequence which is able to bind to a T cell receptor, which T cell receptor recognises an agent as defined in claim 1, and which sequence is able to cause antagonism of a T cell that carries such a T cell receptor.
21. A method of identifying an antagonist of a T cell, which T cell recognises an agent as defined in claim 1, comprising contacting a candidate substance with the T cell and detecting whether the substance causes a decrease in the ability of the T cell to undergo an antigen specific response, the detecting of any such decrease in said ability indicating that the substance is an antagonist.
22. A kit for carrying out a method or use according to any one of claims 1 to 11 comprising an agent as defined in claim 1, 3 or 4 and a means to detect the recognition of the peptide by the T cell.
23. A kit according to claim 22 wherein the means to detect recognition comprises an antibody to IFN- γ .
24. A kit according to claim 23 wherein the antibody is immobilised on a solid support and optionally the kit also comprises a means to detect the

antibody/IFN- γ complex.

25. An agent as defined in claim 1 or an antagonist as defined in claim 15.
 26. A pharmaceutical composition comprising an agent or antagonist as defined in claim 25 and a pharmaceutically acceptable carrier or diluent.
-
27. A composition for tolerising an individual to a gliadin protein to suppress the production of a T cell or antibody response to an agent as defined in claim 1, which composition comprises an agent as defined in claim 1.
 28. A composition for antagonising a T cell response to an agent as defined in claim 1, which composition comprises an antagonist as defined in claim 15.
 29. Use of an agent or antagonist as defined in claim 24 or a wild type sequence as defined in claim 19 to produce an antibody specific to the agent, antagonist or wild type sequence.
 30. Use of a mutation in an epitope of a gliadin protein, which epitope is as defined in claim 1, to decrease the ability of the gliadin protein to cause coeliac disease.
 31. A polynucleotide that comprises a coding sequence that encodes a protein or fragment as defined in claim 19 or 20.
 32. A polynucleotide according to claim 31 that additionally comprises one or more regulatory sequences operably linked to the coding sequence, which regulatory sequences are capable of securing the expression of the coding sequence in a cell.
 33. A polynucleotide according to claim 32 wherein the regulatory sequence(s)

allow expression of the coding sequence in a prokaryotic or mammalian cell.

34. A polynucleotide according to any one of claims 31 to 33 which is a vector or which is in the form of a vector.
35. A cell comprising a polynucleotide as defined in any one of claims 30 to 34 or which has been transformed with such a polynucleotide.
36. A cell according to claim 35 which is a prokaryotic cell or a mammalian cell.
37. A mammal that expresses a T cell receptor as defined in claim 1.
38. Method of identifying a product which is therapeutic for coeliac disease comprising administering a candidate substance to a mammal as defined in claim 37 which has, or which is susceptible to, coeliac disease and determining whether substance prevents or treats coeliac disease in the mammal, the prevention or treatment of coeliac disease indicating that the substance is a therapeutic product.
39. A therapeutic product as identified in the method of claim 38 for use in a method of preventing or treating coeliac disease.
40. A method of diagnosing coeliac disease, or susceptibility to coeliac disease in an individual comprising administering an agent as defined in claim 1 and determining *in vivo* or *in vitro* whether T cells in the individual recognise the agent, recognition of the agent indicating that the individual has or is susceptible to coeliac disease.
41. A method of preventing or treating coeliac disease comprising administering an agent or composition as defined in claim 1 or any one of claims 26 to 28.

42. A method of preventing or treating coeliac disease comprising (a) diagnosing coeliac disease in an individual in a method as defined in claim 1 or claim 40, and (b) administering to an individual diagnosed in (a) as having, or being susceptible to, coeliac disease a therapeutic agent for preventing or treating coeliac disease.
43. A cell according to claim 35 which is a cell of a graminaceous monocotyledonous species.
44. A cell according to claim 43 which is a cell of wheat, maize, oats, rye, rice, barley, triticale, sorghum, or sugar cane.
45. A process for the production of a protein encoded by a coding sequence as defined in claim 31 which process comprises:
- (a) cultivating a cell according to any one of claims 35, 36, 43 or 44 under conditions that allow the expression of the protein; and optionally
 - (b) recovering the expressed protein.
46. A method of obtaining a transgenic plant cell comprising:
- (a) transforming a plant cell with a vector according to claim 34 to give a transgenic plant cell.
47. A method of obtaining a first-generation transgenic plant comprising:
- (b) regenerating a transgenic plant cell transformed with a vector according to claim 34 to give a transgenic plant.
48. A method of obtaining a transgenic plant seed comprising:

- (c) obtaining a transgenic seed from a transgenic plant obtainable by step (b) of claim 47.

49. A method of obtaining a transgenic progeny plant comprising obtaining a second-generation transgenic progeny plant from a first-generation transgenic plant obtainable by a method according to claim 47, and optionally obtaining transgenic plants of one or more further generations from the second-generation progeny plant thus obtained.

50. A method according to claim 49 comprising:

- (d) obtaining a transgenic seed from a first-generation transgenic plant obtainable by the method according to claim 48, then obtaining a second-generation transgenic progeny plant from the transgenic seed;

and/or

- (e) propagating clonally a first-generation transgenic plant obtainable by the method according to claim 47 to give a second-generation progeny plant;

and/or

- (f) crossing a first-generation transgenic plant obtainable by a method according to claim 47 with another plant to give a second-generation progeny plant;

and optionally

- (g) obtaining transgenic progeny plants of one or more further generations from the second-generation progeny plant thus obtained.

51. A transgenic plant cell, plant, plant seed or progeny plant obtainable by a method according to any one of claims 46 to 51.
52. A transgenic plant or plant seed comprising plant cells according to claim 43 or 44.
53. A transgenic plant cell callus comprising plant cells according to claim 43 or 44 obtainable from a transgenic plant cell, first-generation plant, plant seed or progeny as defined in any one of claims 43, 44, or 46 to 50.
54. A plant or callus according to any one of claims claim 51 to 53 which is of a species as defined in claim 43 or 44.
55. A method of obtaining a crop product comprising harvesting a crop product from a plant according to any one of claims 51 to 54 and optionally further processing the harvested product.
56. A method according to claim 55 wherein the plant is a wheat plant and the harvested crop product is grain; optionally further processed into flour or another grain product.
57. A crop product obtainable by a method according to claim 55 or 56.
58. A food that comprises a protein as defined in any claim 19 or 20.
59. A food according to claim 58 in which a protein as defined in claim 19 or 20 is used instead of wild-type gliadin.

ABSTRACT

DIAGNOSTIC AND THERAPEUTIC EPITOPE, AND TRANSGENIC PLANT

A method of diagnosing coeliac disease, or susceptibility to coeliac disease, in an individual comprising:

- (a) contacting a sample from the host with an agent selected from
 - (i) the epitope comprising sequence which is: SEQ ID NO:1 or 2, or an equivalent sequence from a naturally occurring homologue of the gliadin represented by SEQ ID NO:3,
 - (ii) an epitope comprising sequence comprising: SEQ ID NO:1, or an equivalent sequence from a naturally occurring homologue of the gliadin represented by SEQ ID NO:3, which epitope is an isolated oligopeptide derived from a gliadin protein,
 - (iii) an analogue of (i) or (ii) which is capable of being recognised by a T cell receptor that recognises (i) or (ii), which in the case of a peptide analogue is not more than 50 amino acids in length, or
 - (iv) a product comprising two or more agents as defined in (i), (ii) or (iii), and
- (b) determining *in vitro* whether T cells in the sample recognise the agent; recognition by the T cells indicating that the individual has, or is susceptible to, coeliac disease.

Therapeutic compositions which comprise the epitope and gliadin proteins which do not cause coeliac disease are also provided.

a.






Figure 1

THIS PAGE BLANK (USPTO)

b.

Figure 1



THIS PAGE BLANK (USPTO)

a.





Figure 2

THIS PAGE BLANK (USPTO)

b

Figure 2



THIS PAGE BLANK (USPTO)





Figure 3


THIS PAGE BLANK (USPTO)

Figure 4

a.




b.




THIS PAGE BLANK (USPTO)

Figure 5



THIS PAGE BLANK (USPTO)


Figure 6




THIS PAGE BLANK (USPTO)

Figure 7

a.




b.




THIS PAGE BLANK (USPTO)

Figure 8



THIS PAGE BLANK (USPTO)

Figure 9



PCT NO : G800 / 03760

FRASER DAVIS

THIS PAGE BLANK (USPTO)